Formulation of a Healthy Ingredient using Papaya Carotenoid Enriched Vegetable Oils: A Preliminary Study to Improve Carotenoids Stability and Bioaccessibility with O/W Emulsions

CANO M. (1), LARA-ABIA S. (1,2), PEZ-PASCUAL N. (1), LOBO G. (3), WELTI-CHANES J. (2)

1 Instituto de Investigaci en Ciencias de la Alimentaci (CIAL) Madrid Consejo Superior de Investigaciones Cienticas, Madrid, Spain 2 Tecnolico de Monterrey, Escuela de Ingenier y Ciencias, Centro de Biotecnolog FEMSA, Monterrey, NL, Mexico 3 Instituto Canario de Investigaciones Agrarias (ICIA), Tenerife, Spain

The aim of the present work was to improve stability and bioaccessibility of carotenoids from papaya Sweet Mary tissues using carotenoid-enriched vegetable oils (soybean oil, sunflower oil and MCT) to formulate oil-in-water (O/W) emulsions. The use of different concentrations of pectin (1%, 2% and 3%) and homogeneization conditions (time: 2, 3, 4 and 5 min; rpm: 9500, 12000, 14000 and 16000 rpm) and high pressure homogenization (HPH) at 100 MPa for 5 cycles, were evaluated to determine the optimal condition to formulate such emulsions. The standardized in vitro digestion INFOGEST methodology was used to evaluate the bioaccessibility of papaya carotenoids encapsulated by O/W emulsions. Microstructural studies (confocal and optical microscopy) of the obtained emulsions and during their in vitro digestion phases were carried out. Sunflower O/W microemulsions showed smaller mean particle size, higher negative ?-potential and higher viscosity, than soybean O/W microemulsions. Particle size reduction in sunflower microemulsions favoured the bioaccessibility of papaya carotenoids, being lycopene the carotenoid with the higher bioaccessibility (71.4%), followed by ?-carotene (18.4%), ?-cryptoxanthin (15.0%) and ?- cryptoxanthin laurate (7.2%). Similar bioaccessibility was obtained for ?-cryptoxanthin (15.41%), ?-cryptoxanthin laurate (3.54%), ?-carotene (17.89%) and lycopene (64.09%) in soybean emulsions. These results highlight the potential of using carotenoid-enriched vegetable oils to formulate O/W emulsions to enhance carotenoids bioactivity by efficiently preventing their degradation and increasing in vitro bioaccessibility.

This study was funded by project PDI2020-118300RB-C21 of Spanish Ministry of Science and Innovation and Tecnologico de Monterrey (Research Chair Funds C-200)