

## Bio-Converted Spirulina for Nutraceutical Chewing Candy Formulations Rich in L-Glutamic and Gamma-Aminobutyric Acids

**ROCHA J. (1,6,7), BARTKIENE E. (2,3), TOLPEZNIKAITE E. (2), KLUPSAITE D. (2), STARKUTE V. (2,3), BARTKEVICIS V. (4), SKRASTINA A. (4), PAVLENKO R. (4), MOCKUS E. (2), LELE V. (2,3), BATKEVICIUTE G. (2), BUDRIKYTE A. (2), JANULYTE R. (2), JOMANTAITE I. (2), KYBARTAITE A. (2), KNYSTAUTAITE K. (2), VALIONYTE A. (2), RUIBYS R. (5)**

1 Universidade Catolica Portuguesa, CBQF - Centro de Biotecnologia e Quimica Fina Laboratio Associado, Escola Superior de Biotecnologia, Porto, Portugal

2 Institute of Animal Rearing Technologies, Faculty of Animal Sciences, Lithuanian University of Health Sciences, Kaunas, Lithuania

3 Department of Food Safety and Quality, Faculty of Veterinary, Lithuanian University of Health Sciences, Kaunas, Portugal

4 Institute of Food Safety, Animal Health and Environment BIOR, Riga, Latvia

5 Institute of Agricultural and Food Sciences, Agriculture Academy, Vytautas Magnus University, Kaunas, Lithuania

6 LEPABE-Laboratory for Process Engineering, Environment, Biotechnology and Energy, Department of Chemical Engineering (DEQ), Faculty of Engineering, University of Porto (FEUP), Porto, Portugal

7 ALICE Associate Laboratory in Chemical Engineering, Faculty of Engineering, University of Porto, Porto, Portugal

This study aimed at evaluating changes of microalgae *Spirulina* during its fermentation with *Lactiplantibacillus plantarum* No. 122 strain, and further at incorporating *Spirulina* bio-converted for nutraceuticals rich in L-glutamic (L-Glu) and gamma-aminobutyric acids (GABA) into sucrose-free chewing candy (gummy) preparations. Fermented *Spirulina* had higher  $b^*$  (yellowness) coordinates than untreated (non-fermented), and fermentation duration (24 and 48 h) had a statistically significant effect on colour coordinates. The highest contents of L-glutamic and gamma-aminobutyric acids (4062 and 228.6 mg/kg, respectively) were found in 24 and 48 h-fermented *Spirulina*, respectively. Fermentation increased the content of saturated fatty acids and omega-3 in *Spirulina*, while monounsaturated fatty acids and omega-6 were reduced. The addition of fermented *Spirulina* (FSp) significantly affected hardness, decreased lightness and yellowness, and increased the greenness of chewing candies. All chewing candy samples (with xylitol) prepared with 3 and 5 g of FSp and 0.2  $\mu$ L of Citrus paradise essential oil received the highest scores for overall acceptability, and the highest intensity (0.052) of emotion "happy" was elicited by the sample group containing xylitol, agar, ascorbic acid, 3 g of FSp, and 0.1  $\mu$ L of *Mentha spicata* essential oil. As an outcome of this research, one may conclude that fermented *Spirulina* has significant potential as an innovative ingredient in the production of healthier sucrose-free nutraceutical chewing candies.